

MULTIPLY CONJUGATE SYSTEMS CONTAINING DEFORMATIONS OF QUADRICS

ION I. DINCĂ

ABSTRACT. We provide a generalization of Bianchi's triply conjugate systems containing a family of deformations of 2-dimensional quadrics together with its Bäcklund transformation to higher dimensions.

CONTENTS

1. Introduction	1
2. Confocal quadrics in canonical form	2
3. Multiply conjugate systems containing deformations of quadrics	5
3.1. (Isotropic) quadrics without center	6
3.2. Quadrics with center	9
4. The Bäcklund transformation	9
4.1. (Isotropic) quadrics without center	9
4.2. Quadrics with center	10
References	10

1. INTRODUCTION

At the end of the XIXth century Weingarten produced the first examples of triply orthogonal systems (Lamé families) containing a family of *constant Gauß curvature* (CGC) $K = -1$ surfaces. Bianchi extended his work by allowing the (negative) CGC to vary within its family of surfaces and by developing the *Bäcklund* (B) transformation for such triply orthogonal systems. With the development of the theory of deformations of quadrics at the beginning of the XXth century Bianchi introduced triply conjugate systems containing a family of deformations of quadrics together with its B transformation as the natural metric-projective generalization of Weingarten's triply orthogonal systems (see Darboux [9] for triply conjugate systems, Green [10] for projective differential geometry of triple systems of surfaces and Bianchi ([5], Vol 4,(143)&(146)) for triply conjugate systems containing deformations of quadrics).

According to Bianchi such triply conjugate systems must induce on the deformations of quadrics the conjugate system common to the quadric and its deformation, thus generalizing lines of curvature on CGC surfaces (surfaces of triply orthogonal systems cut each other along lines of curvature).

Note that Peterson's 1-dimensional family of deformations of quadrics (see [11]) a-priori does not provide a triply conjugate system, although we have the conjugate system common to the quadric and its deformation property for each deformation.

In what concerns deformations of higher dimensional quadrics the first results are those of Cartan [7] concerning the deformation problem of space forms in space forms; in particular he introduced the minimal co-dimension $(n-1)$ for the deformation problem of n -dimensional space forms in space

Key words and phrases. Bäcklund transformation, Bianchi Permutability Theorem, common conjugate systems, (confocal) quadrics, (discrete) deformations in $\mathbb{C}^{2n-1}$ of quadrics in $\mathbb{C}^{n+1}$, multiply conjugate systems.

Supported by the University of Bucharest.

forms, the exteriorly orthogonal forms tool (naturally appearing from the Gauß equations), their canonical form (given by lines of curvature on the deformation) and the $n(n-1)$ functions of one variable dimensionality of the space of such deformations. These have been extended on one hand (upon a suggestion from S. S. Chern and using a result due to Moore on the Chebyshev coordinates on deformations of $\mathbb{H}^n(\mathbb{R})$ in $\mathbb{R}^{2n-1}$; these are lines of curvature and thus in bijective correspondence with such deformations) to the B transformation of $\mathbb{H}^n$ in $\mathbb{R}^{2n-1}$ in Tenenblat-Terng [12] (Terng also developed the *Bianchi Permutability Theorem* (BPT) for this B transformation in [13]) and on the other hand to deformations in $\mathbb{R}^{2n-1}$ of quadrics in $\mathbb{R}^{n+1}$ or in $\mathbb{R}^n \times (i\mathbb{R})$ with positive definite linear element in Berger, Bryant and Griffiths [1] (again we have the minimal co-dimension $(n-1)$ and the $n(n-1)$ functions of one variable dimensionality of the space of such deformations).

Note also that multiply orthogonal systems are present in the current literature of integrable systems (see for example Terng-Uhlenbeck [14] and its references).

Thus the natural question of completing deformations of higher dimensional quadrics and their B transformation to multiply conjugate systems containing deformations of higher dimensional quadrics and their B transformation arises. Just as the theory of deformation of 2-dimensional quadrics admits discretization via the iteration of the B transformation (moving Möbius configurations in Bianchi's denomination; see Bobenko-Pinkall [6] for discrete deformations of the 2-dimensional pseudo-sphere), a similar approach should give discrete multiply conjugate systems containing deformations of higher dimensional quadrics (note that Weingarten discovered his triply orthogonal systems of CGC -1 surfaces by the iteration of an infinitesimal transformation of CGC -1 surfaces, so he essentially used the discrete version to find the differential version).

According to the principles laid down by Bianchi, if we restrict $(n-1)$ parameters in the multiply conjugate systems to constants so as to obtain a deformation of a quadric, then the remaining n parameters should provide the conjugate system common to the quadric and its deformation (a-priori the quadric may vary with the $(n-1)$ constants, but we restrict our discussion only to deformations of a fixed quadric).

Thus to obtain multiply conjugate systems containing deformations of quadrics one must extend the differential system for deformations of n -dimensional quadrics in $\mathbb{C}^{2n-1}$ via isothermal-conjugate system on the considered quadric and the conjugate system common to the quadric and its deformation according to Bianchi's principles and Cartan's exterior differential calculus.

For more details on the (classical) theory of deformations of (higher dimensional) quadrics we refer the reader to one of our previous notes concerning Bianchi's Bäcklund transformation for higher dimensional quadrics.

All computations are local and assumed to be valid on their open domain of validity without further details; all functions have the assumed order of differentiability and are assumed to be invertible, non-zero, etc when required (for all practical purposes we can assume all functions to be analytic).

2. CONFOCAL QUADRICS IN CANONICAL FORM

Consider the complexified Euclidean space

$$(\mathbb{C}^m, \langle \cdot, \cdot \rangle), \quad \langle x, y \rangle := x^T y, \quad |x|^2 := x^T x, \quad x, y \in \mathbb{C}^m$$

with standard basis $\{e_j\}_{j=1,\dots,m}$, $e_j^T e_k = \delta_{jk}$.

Isotropic (null) vectors are those vectors v of length 0 ($|v|^2 = 0$); since most vectors are not isotropic we shall call a vector simply vector and we shall only emphasize isotropic when the vector is assumed to be isotropic. The same denomination will apply in other settings: for example we call quadric a non-degenerate quadric (a quadric projectively equivalent to the complex unit sphere).

A quadric $x \subset \mathbb{C}^{n+1}$ is given by the quadratic equation $Q(x) := \begin{bmatrix} x \\ 1 \end{bmatrix}^T \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \begin{bmatrix} x \\ 1 \end{bmatrix} = x^T(Ax + 2B) + C = 0$, $A = A^T \in \mathbf{M}_{n+1}(\mathbb{C})$, $B \in \mathbb{C}^{n+1}$, $C \in \mathbb{C}$, $\begin{vmatrix} A & B \\ B^T & C \end{vmatrix} \neq 0$.

A metric classification of all (totally real) quadrics in $\mathbb{C}^{n+1}$ requires the notion of *symmetric Jordan* (SJ) canonical form of a symmetric complex matrix. The symmetric Jordan blocks are: $J_1 := 0 = 0_{1,1} \in \mathbf{M}_1(\mathbb{C})$, $J_2 := f_1 f_1^T \in \mathbf{M}_2(\mathbb{C})$, $J_3 := f_1 e_3^T + e_3 f_1^T \in \mathbf{M}_3(\mathbb{C})$, $J_4 := f_1 \bar{f}_2^T + f_2 \bar{f}_1^T \in \mathbf{M}_4(\mathbb{C})$, $J_5 := f_1 \bar{f}_2^T + f_2 e_5^T + e_5 \bar{f}_2^T + \bar{f}_2 f_1^T \in \mathbf{M}_5(\mathbb{C})$, $J_6 := f_1 \bar{f}_2^T + f_2 \bar{f}_3^T + f_3 \bar{f}_3^T + \bar{f}_3 \bar{f}_2^T + f_2 \bar{f}_1^T \in \mathbf{M}_6(\mathbb{C})$, etc, where $f_j := \frac{e_{2j-1} + i e_{2j}}{\sqrt{2}}$ are the standard isotropic vectors (at least the blocks J_2, J_3 were known to the classical geometers). Any symmetric complex matrix can be brought via conjugation with a complex rotation to the symmetric Jordan canonical form, that is a matrix block decomposition with blocks of the form $a_j I_p + J_p$; totally real quadrics are obtained for eigenvalues a_j of the quadratic part A defining the quadric being real or coming in complex conjugate pairs $a_j, \bar{a}_j$ with subadjacent symmetric Jordan blocks of same dimension p . Just as the usual Jordan block $\sum_{j=1}^p e_j e_{j+1}^T$ is nilpotent with e_{p+1} cyclic vector of order p , J_p is nilpotent with $\bar{f}_1$ cyclic vector of order p , so we can take square roots of SJ matrices without isotropic kernels ($\sqrt{a I_p + J_p} := \sqrt{a} \sum_{j=0}^{p-1} \binom{\frac{1}{2}}{j} a^{-j} J_p^j$, $a \in \mathbb{C}^*$, $\sqrt{a} := \sqrt{r} e^{i\theta}$ for $a = r e^{2i\theta}$, $0 < r$, $-\pi \leq 2\theta < \pi$), two matrices with same SJ decomposition type (that is J_p is replaced with a polynomial in J_p) commute, etc.

The confocal family $\{x_z\}_{z \in \mathbb{C}}$ of a quadric $x_0 \subset \mathbb{C}^{n+1}$ in canonical form (depending on as few constants as possible) is given in the projective space $\mathbb{CP}^{n+1}$ by the equation $Q_z(x_z) := \begin{bmatrix} x_z \\ 1 \end{bmatrix}^T \left(\begin{bmatrix} A & B \\ B^T & C \end{bmatrix}^{-1} - z \begin{bmatrix} I_{n+1} & 0 \\ 0^T & 0 \end{bmatrix} \right)^{-1} \begin{bmatrix} x_z \\ 1 \end{bmatrix} = 0$, where

- $A = A^T \in \mathbf{GL}_{n+1}(\mathbb{C})$ SJ, $B = 0 \in \mathbb{C}^{n+1}$, $C = -1$ for *quadrics with center* (QC),
- $A = A^T \in \mathbf{M}_{n+1}(\mathbb{C})$ SJ, $\ker(A) = \mathbb{C} e_{n+1}$, $B = -e_{n+1}$, $C = 0$ for *quadrics without center* (QWC) and
- $A = A^T \in \mathbf{M}_{n+1}(\mathbb{C})$ SJ, $\ker(A) = \mathbb{C} f_1$, $B = -\bar{f}_1$, $C = 0$ for *isotropic quadrics without center* (IQC).

From the definition one can see that the family of quadrics confocal to x_0 is the adjugate of the pencil generated by the adjugate of x_0 and Cayley's absolute $C(\infty) \subset \mathbb{CP}^n$ in the hyperplane at infinity; since Cayley's absolute encodes the Euclidean structure of $\mathbb{C}^{n+1}$ (it is the set invariant under rigid motions and homotheties of $\mathbb{C}^{n+1} := \mathbb{CP}^{n+1} \setminus \mathbb{CP}^n$) the mixed metric-projective character of the confocal family becomes clear.

For QC $\text{spec}(A)$ is unambiguous (does not change under rigid motions) but for (I)QWC it may change with $(p+1)$ -roots of unity for the block of $\ker(A)$ in A being J_p even under rigid motions $(R, t) \in \mathbf{O}_{n+1}(\mathbb{C}) \ltimes \mathbb{C}^{n+1}$ which preserve the canonical form, so it is unambiguous up to $(p+1)$ -roots of unity.

We have the diagonal Q(W)C respectively for $A = \sum_{j=1}^{n+1} a_j^{-1} e_j e_j^T$, $A = \sum_{j=1}^n a_j^{-1} e_j e_j^T$; the diagonal IQWC come in different flavors, according to the block of f_1 : $A = J_p + \sum_{j=p+1}^{n+1} a_j^{-1} e_j e_j^T$; in particular if $A = J_{n+1}$, then $\text{spec}(A) = \{0\}$ is unambiguous. General quadrics are those for which all eigenvalues have geometric multiplicity 1; equivalently each eigenvalue has an only corresponding SJ block; in this case the quadric also admits elliptic coordinates.

There are continuous groups of symmetries which preserve the SJ canonical form for more than one SJ block corresponding to an eigenvalue, so from a metric point of view a metric classification according to the elliptic coordinates and continuous symmetries may be a better one.

With $R_z := I_{n+1} - zA$, $z \in \mathbb{C} \setminus \text{spec}(A)^{-1}$ the family of quadrics $\{x_z\}_z$ confocal to x_0 is given by $Q_z(x_z) = x_z^T A R_z^{-1} x_z + 2(R_z^{-1} B)^T x_z + C + z B^T R_z^{-1} B = 0$. For $z \in \text{spec}(A)^{-1}$ we obtain singular confocal quadrics; those with z^{-1} having geometric multiplicity 1 admit a singular set which is an $(n-1)$ -dimensional quadric projectively equivalent to $C(\infty)$, so they will play an important rôle in the discussion of homographies $H \in \mathbf{PGL}_{n+1}(\mathbb{C})$ taking a confocal family into another one, since $H^{-1}(C(\infty))$, $C(\infty)$ respectively $C(\infty)$, $H(C(\infty))$ will suffice to determine each confocal family.

The Ivory affinity is an affine correspondence between confocal quadrics and having good metric properties (it may be the reason why Bianchi calls it *affinity* in more than one language): it is given by $x_z = \sqrt{R_z} x_0 + C(z)$, $C(z) := -(\frac{1}{2} \int_0^z (\sqrt{R_w})^{-1} dw) B$. Note that $C(z) = 0$ for QC,

$= \frac{z}{2}e_{n+1}$ for QWC; for IQWC it is the Taylor series of $\frac{1}{2}\int_0^z(\sqrt{1-w})^{-1}dw$ at $z=0$ with each monomial z^{k+1} replaced by $z^{k+1}J_p^k\bar{f}_1$, where J_p is the block of f_1 in A and thus a polynomial of degree p in z . Note $AC(z) + (I_{n+1} - \sqrt{R_z})B = 0 = (I_{n+1} + \sqrt{R_z})C(z) + zB$. Applying d to $Q_z(x_z) = 0$ we get $dx_z^T R_z^{-1}(Ax_z + B) = 0$, so the unit normal N_z is proportional to $\hat{N}_z := -2\partial_z x_z$. If $\mathbb{C}^{n+1} \ni x \in x_{z_1}, x_{z_2}$, then $\hat{N}_{z_j} = R_{z_j}^{-1}(Ax + B)$; using $R_z^{-1} - I_{n+1} = zAR_z^{-1}$, $z_1R_{z_1}^{-1} - z_2R_{z_2}^{-1} = (z_1 - z_2)R_{z_1}^{-1}R_{z_2}^{-1}$ we get $0 = Q_{z_1}(x) - Q_{z_2}(x) = (z_1 - z_2)\hat{N}_{z_1}^T\hat{N}_{z_2}$, so two confocal quadrics cut each other orthogonally (Lamé). For general quadrics the polynomial equation $Q_z(x) = 0$ has degree $n+1$ in z and it has multiple roots iff $0 = \partial_z Q_z(x) = |\hat{N}_z|^2$; thus outside the locus of isotropic normals elliptic coordinates (given by the roots $z_1, \dots, z_{n+1}$ of the said equation) give a parametrization of $\mathbb{C}^{n+1}$ suited to confocal quadrics.

We have now some classical metric properties of the Ivory affinity: with $x_0^0, x_0^1 \in x_0$, $V_0^1 := x_z^1 - x_0^0$, etc the Ivory Theorem (preservation of length of segments between confocal quadrics) becomes $|V_0^1|^2 = |x_0^0 + x_0^1 - C(z)|^2 - 2(x_0^0)^T(I_{n+1} + \sqrt{R_z})x_0^1 + zC = |V_0^1|^2$; the preservation of lengths of rulings: $w_0^T Aw_0 = w_0^T \hat{N}_0 = 0$, $w_z = \sqrt{R_z}w_0 \Rightarrow w_z^T w_z = |w_0|^2 - zw_0^T Aw_0 = |w_0|^2$; the symmetry of the *tangency configuration* (TC): $(V_0^1)^T \hat{N}_0^0 = (x_0^0)^T A \sqrt{R_z} x_0^1 - B^T (x_0^0 + x_z^1 - C(z)) + C = (V_0^1)^T \hat{N}_0^1$; the preservation of angles between segments and rulings: $(V_0^1)^T w_0^0 + (V_0^1)^T w_z^0 = -z(\hat{N}_0^0)^T w_0^0 = 0$; the preservation of angles between rulings: $(w_0^0)^T w_z^1 = (w_0^0)^T \sqrt{R_z} w_0^1 = (w_z^0)^T w_0^1$; the preservation of angles between polar rulings: $(w_0^0)^T A \hat{w}_0^0 = 0 \Rightarrow (w_z^0)^T \hat{w}_z^0 = (w_0^0)^T \hat{w}_0^0 - z(w_0^0)^T A \hat{w}_0^0 = (w_0^0)^T \hat{w}_0^0$.

All complex quadrics are affine equivalent to either the unit sphere $X \subset \mathbb{C}^{n+1}$, $|X|^2 = 1$ or to the equilateral paraboloid $Z \subset \mathbb{C}^{n+1}$, $Z^T(I_{1,n}Z - 2e_{n+1}) = 0$, so a parametrization with regard to these two quadrics is in order: $x_0 = (\sqrt{A})^{-1}X$ for QC, $x_0 = (\sqrt{A + e_{n+1}e_{n+1}^T})^{-1}Z$ for QWC (for this reason from a canonical metric point of view (that is we are interested in a simplest form of $|dx_0|^2$) we should rather require that A^{-1} or $(A + e_{n+1}e_{n+1}^T)^{-1}$ is SJ).

For IQWC such a parametrization fails because of the isotropic $\ker(A)$; however the computations between confocal quadrics involving the Ivory affinity reveal a natural parametrization of IQWC which is again an affine transformation of Z .

Consider a canonical IQWC $x_0^T(Ax_0 - 2\bar{f}_1) = 0$, $\ker(A) = \mathbb{C}f_1$, $A = J_p \oplus \dots$ SJ. We are looking for a linear map $L \in \mathbf{GL}_{n+1}(\mathbb{C})$ such that $x_0 = LZ$, equivalently $L^T AL = e^{2a}I_{1,n}$, $I_{1,n} := I_{n+1} - e_{n+1}e_{n+1}^T$, $L^T \bar{f}_1 = e^{2a}e_{n+1}$. Replacing L with $L(e^{-a}I_{1,n} + e^{-2a}e_{n+1}e_{n+1}^T)$ we can make $a = 0$. Thus $Le_{n+1} = f_1$, $L^T(A + \bar{f}_1\bar{f}_1^T)L = I_{n+1}$, so $L^{-1} = R^T\sqrt{A + \bar{f}_1\bar{f}_1^T}$, $R^T R = I_{n+1}$ with $Re_{n+1} = \sqrt{A + \bar{f}_1\bar{f}_1^T}f_1$ (note that Re_{n+1} has, as required, length 1). Once $R \in \mathbf{O}_{n+1}(\mathbb{C})$ with the above property is found, L thus defined satisfies $L^T \bar{f}_1 = e_{n+1}$ and thus $L^T AL = I_{1,n}$. L with the above properties is unique up to rotations fixing e_{n+1} in its domain and a canonical choice of R reveals itself from a SJ canonical form when doing computations on confocal quadrics. We have $LL^T = (A + \bar{f}_1\bar{f}_1^T)^{-1}$, $I_{1,n}L^{-1}\sqrt{R_z}L = I_{1,n}L^{-1}\sqrt{R_z}LI_{1,n} = L^{-1}\sqrt{R_z}L - e_{n+1}\bar{f}_1\sqrt{R_z}L = L^T A \sqrt{R_z}L = I_{1,n}\sqrt{I_{n+1} - zL^T A^2 L} =: I_{1,n}\sqrt{R'_z}$, $A' := L^T A^2 L$, $\ker(A') = \mathbb{C}e_{n+1} \oplus \mathbb{C}L^{-1}(A + \bar{f}_1\bar{f}_1^T)^{-1}f_1 = \mathbb{C}e_{n+1} \oplus \mathbb{C}L^T f_1$; choose R which makes A' SJ. Note that we can take for QWC $L := (\sqrt{A + e_{n+1}e_{n+1}^T})^{-1}$, $A' := A$, $\ker(A') = \mathbb{C}e_{n+1}$, so IQWC can be regarded as metrically degenerated QWC. Note that $e_{n+1}^T L^{-1}\sqrt{R_z}L = (-I_{1,n}L^{-1}C(z) + e_{n+1})^T$; this can be confirmed analytically by differentiating with respect to z and using $(L^T)^{-1} = AL - Be_{n+1}^T$ and will imply the symmetry of the TC, but since we have already proved the symmetry of the TC, we can use this to imply the previous. Thus $L^{-1}x_z = L^{-1}\sqrt{R_z}LZ + L^{-1}C(z) = I_{1,n}\sqrt{R'_z}Z + e_{n+1}(-I_{1,n}L^{-1}C(z) + e_{n+1})^T Z + L^{-1}C(z)$, $(x_z^1 - x_0^0)^T \hat{N}_0^0 = (L^{-1}x_z^1 - Z_0)^T(I_{1,n}Z_0 - e_{n+1}) = Z_0^T I_{1,n}\sqrt{R'_z}Z_1 + (Z_0 + Z_1)^T(I_{1,n}L^{-1}C(z) - e_{n+1}) - e_{n+1}^T L^{-1}C(z)$. Note that for IQWC $|I_{1,n}L^{-1}C(z)|^2 = 2e_{n+1}^T L^{-1}C(z)$, so $L^{-1}C(z)$ lies itself on Z (also in this case since $\bar{f}_1^T J_p^k \bar{f}_1 = \delta_{k,p-1}$ we have $e_{n+1}^T L^{-1}C(z) = \bar{f}_1^T C(z) = \binom{-\frac{1}{2}}{p-1} \frac{(-z)^p}{-2p}$, so e_{n+1}^T picks up the highest power of z in $L^{-1}C(z)$). To see this we need $0 = |L^{-1}C(z) - \bar{f}_1^T C(z)e_{n+1}|^2 - 2\bar{f}_1^T C(z) = C(z)^T(LL^T)^{-1}C(z) - (\bar{f}_1^T C(z))^2 - 2\bar{f}_1^T C(z) = C(z)^T AC(z) - 2\bar{f}_1^T C(z)$; using $AC(z) = (I_{n+1} - \sqrt{R_z})\bar{f}_1$ and $(I_{n+1} + \sqrt{R_z})C(z) = z\bar{f}_1$ it is satisfied.

Note also that as needed later we have $(L^T L)^{-1} = A' - I_{1,n} L^{-1} B e_{n+1}^T - e_{n+1} (I_{1,n} L^{-1} B)^T + |B|^2 e_{n+1} e_{n+1}^T$, $|\hat{N}_0|^2 = |(L^T)^{-1} (I_{1,n} Z - e_{n+1})|^2 = Z^T A' Z + 2Z^T I_{1,n} L^{-1} B + |B|^2$, $(I_{n+1} + \sqrt{R'_z}) I_{1,n} L^{-1} C(z) = I_{1,n} L^{-1} (I_{n+1} + \sqrt{R_z}) C(z) = -z I_{1,n} L^{-1} B$.

3. MULTIPLY CONJUGATE SYSTEMS CONTAINING DEFORMATIONS OF QUADRICS

We assume all subspaces in discussion to be non-isotropic (the Euclidean product induced on them by the one on $\mathbb{C}^m$ is non-degenerate; this assures the existence of orthonormal normal frames and thus the discussion of sub-manifolds via the *Gauß-Weingarten* (GW) and *Gauß-Codazzi-Mainardi(-Peterson)-Ricci* (G-CMP-R) equations).

For deformations $x \subset \mathbb{C}^{2n-1}$ of $x_0 \subset \mathbb{C}^{n+1}$ (that is $|dx|^2 = |dx_0|^2$) with common conjugate system $(u^1, \dots, u^n)$ and non-degenerate joined second fundamental forms (that is $[d^2 x_0^T N_0 \quad d^2 x^T N]$ is a symmetric quadratic $\mathbb{C}^n$ -valued form which contains only $(du^j)^2$ terms for N_0 unit normal field of x_0 and $N = [N_1 \dots N_{n-1}]$ orthonormal normal frame of x and the dimension n cannot be lowered in an open dense set) the linear element must satisfy the condition

$$\Gamma_{jk}^l = 0, \quad j, k, l \text{ distinct}$$

and such deformations $x \subset \mathbb{C}^{2n-1}$ are in bijective correspondence with solutions $\{\mathbf{a}_j\}_{j=1, \dots, n} \subset \mathbf{C}^*$ of the differential system

$$(3.1) \quad (\log \mathbf{a}_j)_k = \Gamma_{jk}^j, \quad j \neq k, \quad \sum_{j=1}^n \frac{(h_j^0)^2}{\mathbf{a}_j^2} + 1 = 0,$$

where $N_0^T d^2 x_0 =: \sum_{j=1}^n h_j^0 (du^j)^2$ is the second fundamental form of x_0 (we shall use Latin indices $j, k, l, \dots$ including to differentiate respectively with $u^j, u^k, u^l, \dots$ when clear from the context; also we shall preserve the classical notation d^2 for the symmetric (tensorial) second derivative and we shall use $d\wedge$ for the exterior (antisymmetric) derivative; thus $d \wedge d = 0$).

Once a solution of this system is known, one finds the second fundamental form of x (complete the row $[i \frac{h_1^0}{\mathbf{a}_1} \dots i \frac{h_n^0}{\mathbf{a}_n}]_j$ as the first row in an orthogonal $R \subset \mathbf{O}_n(\mathbb{C})$, delete it, multiply the column j of the $\mathbf{M}_{n-1,n}(\mathbb{C})$ obtained matrix respectively with $\mathbf{a}_j$ and take the k -th row to obtain the second fundamental form of x in the N_k direction) and then one finds x by the integration of a Ricatti equation and quadratures (the Gauß-Bonnet(-Peterson) Theorem).

By the argument of Cartan's reduction of exteriorly orthogonal forms to the canonical form such coordinates $\{u^j\}_j$ exist for real deformations $\subset \mathbb{R}^{2n-1}$ of imaginary quadrics $\subset \mathbb{R}^n \times i\mathbb{R}$ of negative curvature (for such cases also all computations in the deformation problem will be real; see Berger, Bryant, Griffiths [1]). However, since we have completely integrable differential systems (systems in involution) for the deformation problem for quadrics, the dimensionality of the space of deformations of quadrics remains the same (namely solution depending on $n(n-1)$ functions of one variable) in the complex setting also (the Cartan characters remain the same).

To obtain multiply conjugate systems we need to extend x with the independent variables $u^{n+1}, \dots, u^{2n-1}$.

An m -dimensional region

$$x = x(u^1, \dots, u^m) \subseteq \mathbb{C}^m, \quad du^1 \wedge \dots \wedge du^m \neq 0$$

gives a multiply conjugate system iff

$$(3.2) \quad x_{jk} = (\log \mathbf{a}_j)_k x_j + (\log \mathbf{a}_k)_j x_k, \quad \mathbf{a}_j \in \mathbb{C}^*, \quad j, k = 1, \dots, m, \quad j \neq k$$

with the compatibility condition

$$(3.3) \quad (\mathbf{a}_j)_{kl} = (\log \mathbf{a}_k)_l (\mathbf{a}_j)_k + (\log \mathbf{a}_l)_k (\mathbf{a}_j)_l, \quad j, k, l = 1, \dots, m \text{ distinct.}$$

Note that (3.3) is the condition that the Riemann symbols R_{jklp} of the linear element $\sum_{j=1}^m \mathbf{a}_j^2 (du^j)^2$ with at least three of j, k, l, p distinct are 0, so multiply conjugate systems in $\mathbb{C}^m$ are put in correspondence with such linear elements (normal curved spaces in Bianchi's denomination).

For the specific computations of deformations of quadrics we shall use the convention $\mathbb{C}^n \subset \mathbb{C}^{n+1}$ with 0 on the $(n+1)^{\text{th}}$ component; thus for example we can multiply $(n+1, n+1)$ -matrices with n -column vectors and similarly one can extend (n, n) matrices to $(n+1, n+1)$ matrices with zeroes on the last column and row. The converse is also valid: an $(n+1, n+1)$ matrix with zeroes on the last column and row (or multiplied on the left with an n -row vector and on the right with an n -column vector) will be considered as an (n, n) -matrix.

3.1. (Isotropic) quadrics without center.

With $V := \sum_{k=1}^n v^k e_k = [v^1 \dots v^n]^T$ consider the complex equilateral paraboloid $Z = Z(v^1, \dots, v^n) = V + \frac{|V|^2}{2} e_{n+1}$.

We have the (I)QWC $x_0 := LZ$, $L \in \mathbf{GL}_{n+1}(\mathbb{C})$ (recall $L := (\sqrt{A + e_{n+1} e_{n+1}^T})^{-1}$, $\ker(A) = \mathbb{C} e_{n+1}$, A SJ, $B = -e_{n+1}$ for QWC and $Le_{n+1} = f_1$, $L^T(A + \bar{f}_1 \bar{f}_1^T)L = I_{n+1}$, $A' := L^T A^2 L$ SJ for $\ker(A) = \mathbb{C} f_1$, A SJ, $B = -\bar{f}_1$ in the case of IQWC) with linear element, unit normal, second fundamental form and Christoffel symbols $|dx_0|^2 = dV^T L^T L dV + (V^T dV)^2 |Le_{n+1}|^2 + 2(Le_{n+1})^T L dV (V^T dV)$, $N_0 = \frac{(L^T)^{-1}V+B}{\sqrt{H}}$, $N_0^T d^2 x_0 = -\frac{|dV|^2}{\sqrt{H}}$, $H := |(L^T)^{-1}V + B|^2 = V^T A' V + 2V^T L^{-1}B + |B|^2$, $\tilde{\Gamma}_{jk}^l = 0$, $j \neq k$, $\tilde{\Gamma}_{jj}^k = \frac{\partial \log \sqrt{H}}{\partial v^k}$.

Because $(v^1, \dots, v^n)$ are isothermal-conjugate and $(u^1, \dots, u^n)$ are conjugate on x_0 , the Jacobian $\frac{\partial(v^1, \dots, v^n)}{\partial(u^1, \dots, u^n)}$ has orthogonal columns, so with $\lambda_j := |\frac{\partial V}{\partial u^j}| \neq 0$, $\Lambda := [\lambda_1 \dots \lambda_n]^T$, $\delta' := \text{diag}[du^1 \dots du^n]$, $d'f := \sum_{j=1}^n f_j du^j$ we have $d'V = R\delta'\Lambda$, $R \in \mathbf{O}_n(\mathbb{C})$. Multiplying the formula for the change of Christoffel symbols $\frac{\partial v^c}{\partial u^l} \Gamma_{jk}^l = \frac{\partial^2 v^c}{\partial u^j \partial u^k} + \frac{\partial v^a}{\partial u^j} \frac{\partial v^b}{\partial u^k} \tilde{\Gamma}_{ab}^c = \frac{\partial^2 v^c}{\partial u^j \partial u^k} + \lambda_j^2 \delta_{jk} \frac{\partial \log \sqrt{H}}{\partial v^c}$ on the left with $\frac{\partial v^c}{\partial u^p}$ and summing after c we obtain $\Gamma_{jk}^p = \lambda_p^{-2} (\sum_c \frac{\partial v^c}{\partial u^p} \frac{\partial^2 v^c}{\partial u^j \partial u^k} + \lambda_j^2 \delta_{jk} (\log \sqrt{H})_p) = \delta_{pk} (\log \lambda_k)_j + \delta_{jk} \frac{\lambda_j^2}{\lambda_p^2} (\log \frac{\sqrt{H}}{\lambda_j})_p + \delta_{pj} (\log \lambda_j)_k$, so $\Gamma_{jk}^j = (\log \lambda_j)_k$, $\Gamma_{jj}^k = \frac{\lambda_j^2}{\lambda_k^2} (\log \frac{\sqrt{H}}{\lambda_j})_k$, $j \neq k$, $\Gamma_{jj}^j = (\log(\lambda_j \sqrt{H}))_j$. We have $h_j^0 = -\frac{\lambda_j^2}{\sqrt{H}}$; since $(\log \lambda_j)_k = \Gamma_{jk}^j = (\log \mathbf{a}_j)_k$, $j \neq k$ we get $\lambda_j = \phi_j(u^j) \mathbf{a}_j$; after a change of the u^j variable into itself we can make $\lambda_j = \mathbf{a}_j$, $j = 1, \dots, n$, so from (3.1) $|\Lambda|^2 = -H$ and $\Lambda^T d'\Lambda = -d'V^T (A'V + L^{-1}B) = -\Lambda^T \delta' R^T (A'V + L^{-1}B)$.

Imposing the compatibility condition $R^T d'\Lambda$ on $d'V = R\delta'\Lambda$ we get $R^T d'R \wedge \delta'\Lambda - \delta' \wedge d'\Lambda = 0$, or $(\lambda_j)_k = e_j^T R^T R_j e_k \lambda_k$, $j \neq k$, $e_l^T \frac{R^T R_j}{\lambda_j} e_k = e_l^T \frac{R^T R_k}{\lambda_k} e_j$, j, k, l distinct. Now by the standard Cartan trick $-e_k^T \frac{R^T R_l}{\lambda_l} e_j = -e_k^T \frac{R^T R_j}{\lambda_j} e_l = e_l^T \frac{R^T R_j}{\lambda_j} e_k = e_l^T \frac{R^T R_k}{\lambda_k} e_j = -e_j^T \frac{R^T R_k}{\lambda_k} e_l = -e_j^T \frac{R^T R_l}{\lambda_l} e_k = e_k^T \frac{R^T R_l}{\lambda_l} e_j$ for j, k, l distinct, so $e_j^T R^T R_k e_l = 0$ for j, k, l distinct. Keeping account of the prime integral property $\Lambda^T [d'\Lambda + \delta' R^T (A'V + L^{-1}B)] = 0$ and with $\omega' := \sum_{j=1}^n (e_j e_j^T R^T R_j \delta' + \delta' R^T R_j e_j e_j^T) = -\omega'^T$ we have $d'\Lambda = \omega'\Lambda - \delta' R^T (A'V + L^{-1}B)$ and $d' \wedge d'\Lambda = 0$ becomes the differential system

$$(3.4) \quad d' \wedge \omega' = \omega' \wedge \omega' - \delta' R^T A' R \wedge \delta', \quad \omega' \wedge \delta' = \delta' \wedge R^T d'R, \quad R \in \mathbf{O}_n(\mathbb{C})$$

in involution (that is no further conditions appear if one imposes $d' \wedge$ conditions and one uses the equations of the system itself) as the compatibility condition for the completely integrable linear differential system

$$(3.5) \quad d'V = R\delta'\Lambda, \quad d'\Lambda = \omega'\Lambda - \delta' R^T (A'V + L^{-1}B), \quad \Lambda^T \Lambda = -(V^T A'V + 2V^T L^{-1}B + |B|^2).$$

Note that for QWC once we know a solution $R \in \mathbf{O}_n(\mathbb{C})$ of (3.4), a solution V , Λ of (3.5) and certain linearly independent solutions V_j , Λ_j , $j = 1, \dots, 2n-1$ of the (homogeneous) differential part of (3.5), then we can find up to rigid motions the space realization of the deformation $x \subset \mathbb{C}^{2n-1}$ of x_0 up to quadratures and without the use of the Gauß-Bonnet(-Peterson) Theorem (this is again due to Bianchi for $n = 2$).

Let V_j , Λ_j , $j = 1, \dots, 2n-1$ be certain linearly independent solutions of the (homogeneous) differential part of (3.5) and V , Λ solution of (3.5) such that $x \subset \mathbb{C}^{2n-1}$ given up to quadratures

by

$$(3.6) \quad d'x := [V_1 \ \dots \ V_{2n-1}]^T d'V$$

is a deformation of x_0 with non-degenerate joined second fundamental forms. Note $d' \wedge d'x = 0$; since $|d'x|^2 = |d'x_0|^2$, $d'x_0 = (L + e_{n+1}V^T)d'V$ we get $\sum_{j=1}^{2n-1} V_j V_j^T - VV^T = I_{1,n}L^TL$; applying d' we get $\sum_{j=1}^{2n-1} V_j \Lambda_j^T - V\Lambda^T = 0$ (use $\delta'M = N\delta'$, $M, N \subset \mathbf{M}_n(\mathbb{C}) \Leftrightarrow M = N = \text{diag}$); applying d' again we get $\sum_{j=1}^{2n-1} \Lambda_j \Lambda_j^T - \Lambda\Lambda^T = I_n$.

With $\mathcal{V} := [V_1 \ \dots \ V_{2n-1} \ iV]$, $\mathcal{L} := [\Lambda_1 \ \dots \ \Lambda_{2n-1} \ i\Lambda]$ these can be written as

$$(3.7) \quad \begin{bmatrix} \mathcal{V} \\ \mathcal{L} \end{bmatrix} [\mathcal{V}^T \ \mathcal{L}^T] = \begin{bmatrix} I_{1,n}L^TL & 0 \\ 0 & I_n \end{bmatrix}.$$

We also have the prime integral property $[\mathcal{V}^T \ \mathcal{L}^T] \begin{bmatrix} A' & 0 \\ 0 & I_n \end{bmatrix} \begin{bmatrix} \mathcal{V} \\ \mathcal{L} \end{bmatrix} = \begin{bmatrix} \mathcal{C} & ic \\ ic^T & 1 \end{bmatrix}$, $\mathcal{C} = \mathcal{C}^T \in \mathbf{M}_{2n-1}(\mathbb{C})$, $c \in \mathbb{C}^{2n-1}$; multiplying it on the left with $\begin{bmatrix} \mathcal{V} \\ \mathcal{L} \end{bmatrix}$ and using (3.7) we get $\mathcal{C} = I_{2n-1}$, $c = 0$; thus $[\mathcal{V}^T \ \mathcal{L}^T]$

is determined modulo a multiplication on the left with a rotation $\begin{bmatrix} \mathcal{R} & 0 \\ 0 & 1 \end{bmatrix} \in \mathbf{O}_{2n}(\mathbb{C})$ by

$$(3.8) \quad [\mathcal{V}^T \ \mathcal{L}^T] \begin{bmatrix} A' & 0 \\ 0 & I_n \end{bmatrix} \begin{bmatrix} \mathcal{V} \\ \mathcal{L} \end{bmatrix} = I_{2n}.$$

Thus (3.8) is equivalent to (3.7) and to $[\mathcal{V}^T L^{-1} \ \mathcal{L}^T] \subset \mathbf{O}_{2n}(\mathbb{C})$.

For the non-degenerate joined second fundamental forms property we need to prove that there is no vector field $N \subset \mathbb{C}^{2n-1}$ along x such that $N^T d'x = 0$, $N^T d'^2x = |d'V|^2 = \Lambda^T \delta'^2 \Lambda$, that is the linear system $\begin{bmatrix} \mathcal{V} \\ \mathcal{L} \end{bmatrix} \begin{bmatrix} N \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ \Lambda \end{bmatrix}$ is inconsistent; using (3.8) this becomes $\begin{bmatrix} N \\ 0 \end{bmatrix} = \mathcal{L}^T \Lambda$, which is indeed inconsistent because $|\Lambda|^2 \neq 0$.

Note that since for IQWC $\ker A' = \mathbb{C}e_{n+1} \oplus \mathbb{C}L^T f_1$, $\text{coker } A' = \mathbb{C}e_{n+1} \oplus \mathbb{C}L^{-1}B$ for such quadrics we cannot derive from the equivalent of (3.8) for IQWC the full information about the equivalent of (3.7) for IQWC. This is to be expected, since a fundamental set of solutions of the homogeneous differential part of (3.5) contains $2n$ linearly independent solutions; the obvious solution $[f_1^T L \ 0]^T$ is isotropic and perpendicular on all the others *with respect to* (wrt) the bilinear form given by the left hand side of (3.8); by the Gramm-Schmidt orthogonalization process one can complete this solution with $2n-1$ other solutions V_j , Λ_j , $j = 1, \dots, 2n-1$ orthonormal wrt the same bilinear form. This isotropic solution is added in the $2n-1$ other with yet undetermined constant coefficients $V_j^T L^{-1}B$. The part of the equivalent of (3.7) for IQWC that cannot be inferred from the equivalent of (3.8) for IQWC is $\begin{bmatrix} V \\ \Lambda \end{bmatrix} = -\sum_{j=1}^{2n-1} \begin{bmatrix} V_j \\ \Lambda_j \end{bmatrix} V_j^T L^{-1}B + \begin{bmatrix} L^T f_1 \\ 0 \end{bmatrix} V^T L^{-1}B$, but this requires $V^T L^{-1}B$ to be constant, so V , Λ has to be solution of the homogeneous differential part of (3.5), a contradiction, so V_j , Λ_j , $j = 1, \dots, 2n-1$ cannot be solutions of the homogeneous differential part of (3.5).

We now extend the independent variables $u^1, \dots, u^n$ with independent variables $u^{n+1}, \dots, u^{2n-1}$, $du^1 \wedge \dots \wedge du^{2n-1} \neq 0$, $d = d' + d''$, $d'f = \sum_{j=1}^n f_j du^j$, $d''f = \sum_{j=1}^{n-1} f_{j+n} du^{j+n}$; thus $d \wedge d = 0$ becomes $d' \wedge d' = 0$, $d' \wedge d'' + d'' \wedge d' = 0$, $d'' \wedge d'' = 0$.

With $\delta'' := \text{diag}[du^{n+1} \ \dots \ du^{2n-1} \ 0]$ we now have $d''\Lambda = M\delta''V$, $M \subset \mathbf{M}_n(\mathbb{C})$, $Me_n = 0$ (this choice of M is due to Bianchi for $n = 2$ and its reason will appear immediately).

Imposing the compatibility condition $d' \wedge d''\Lambda + d'' \wedge d'\Lambda = 0$ we obtain $[d' \wedge (M\delta'') - \omega' \wedge M\delta'']V + (d'' \wedge \omega' - M\delta'' \wedge R\delta')\Lambda + \delta'R^T \wedge (A'd''V - d''RR^T(A'V + L^{-1}B)) = 0$, from where we obtain by applying $\delta'\Lambda$: $\delta' \wedge [d' \wedge (M\delta'') - \omega' \wedge M\delta''] = 0$, $\delta' \wedge (d'' \wedge \omega' - M\delta'' \wedge R\delta') = 0$, so $d' \wedge (M\delta'') = \omega' \wedge M\delta'' + \delta'R^T \wedge N\delta''$, $N \subset \mathbf{M}_n(\mathbb{C})$, $Ne_n = 0$, $d'' \wedge \omega' = M\delta'' \wedge R\delta' + \delta'R^T \wedge \dots$. Since $\omega'^T = -\omega'$ we have $d'' \wedge \omega' = M\delta'' \wedge R\delta' + \delta'R^T \wedge \delta''M^T$ (in particular we get M from R and its derivatives; this is the reason for the choice of M), so $A'd''V = (d''RR^T A' - N\delta'')V - \delta''M^T \Lambda + d''RR^T L^{-1}B$. Imposing the compatibility condition $d' \wedge A'd''V + d'' \wedge A'd'V = 0$ and collecting the coefficient of $R\delta'\Lambda$ we get

$d''RR^TA' - N\delta'' = A'd''RR^T + \delta''N^T$ (in particular we get N from R and its derivatives). Note that since for IQWC $\ker A' = \mathbb{C}e_{n+1} \oplus \mathbb{C}L^T f_1$, $\text{coker } A' = \mathbb{C}e_{n+1} \oplus \mathbb{C}L^{-1}B$ for such quadrics we cannot derive the full information about $d''V$ from $A'd''V$ only: from the prime integral property $d''\Lambda^T\Lambda = -d''V^T(A'V + L^{-1}B)$ and using $V^T(d''RR^TA' - N\delta'')V = 0$ we have $(L^{-1}B)^T(d''V - d''RR^TV) = 0$. Now using $A'(d''V - d''RR^TV) = [\mathcal{A}^{-1} + e_{n+1}(L^{-1}B)^T](d''V - d''RR^TV)$, $\mathcal{A} := L^TL$ we can finally extend the completely integrable linear differential system (3.5) to the completely integrable linear differential system

$$(3.9) \quad dV = R\delta'\Lambda + (d''RR^T + \mathcal{A}\delta''N^T)V - \mathcal{A}\delta''M^T\Lambda + \mathcal{A}d''RR^TL^{-1}B, \\ d\Lambda = \omega'\Lambda - \delta'R^T(A'V + L^{-1}B) + M\delta''V, \quad \Lambda^T\Lambda = -(V^TA'V + 2V^TL^{-1}B + |B|^2)$$

with the extended compatibility (and algebraic) conditions

$$(3.10) \quad d' \wedge \omega' = \omega' \wedge \omega' - \delta'R^TA'R \wedge \delta', \quad \omega' \wedge \delta' = \delta' \wedge R^Td'R, \quad d'' \wedge \omega' = M\delta'' \wedge R\delta' + \delta'R^T \wedge \delta''M^T \\ (\Leftrightarrow R d'' \wedge (R^T d'R) = -d' \wedge (d''RR^T)R \quad d' \wedge (d''RR^T) = \delta''M^T \wedge \delta'R^T + R\delta' \wedge M\delta''), \\ d' \wedge (M\delta'') = \omega' \wedge M\delta'' + \delta'R^T \wedge N\delta'', \quad d' \wedge (N\delta'') = R\delta' \wedge M\delta''A' - A'R\delta' \wedge M\delta'', \\ d'' \wedge (M\delta'') = M\delta'' \wedge (d''RR^T + \mathcal{A}\delta''N^T), \quad d'' \wedge (N\delta'') = \delta''M^T \wedge M\delta'' + \\ N\delta'' \wedge (d''RR^T + \mathcal{A}\delta''N^T) + d''RR^T \wedge N\delta'', \quad A'd''RR^T + \delta''N^T = d''RR^TA' - N\delta'', \\ M\delta''\mathcal{A} \wedge \delta''M^T = 0, \quad e_{n+1}^T\mathcal{A}\delta''M^T = e_{n+1}^T\mathcal{A}\delta''N^T = 0, \quad e_{n+1}^T\mathcal{A}d''RR^TL^{-1}B = 0, \\ M\delta''L^{-1}B = M\delta''\mathcal{A} \wedge d''RR^TL^{-1}B = N\delta''\mathcal{A} \wedge d''RR^TL^{-1}B = 0, \quad R \subset \mathbf{O}_n(\mathbb{C}).$$

If we impose $d' \wedge$ and $d'' \wedge$ on (3.10), use $d' \wedge d' = 0$, $d' \wedge d'' + d'' \wedge d' = 0$, $d'' \wedge d'' = 0$ and the equations of the system itself, then we get the algebraic conditions

$$(3.11) \quad M\delta''\mathcal{A} \wedge \delta''N^T \wedge R\delta' + \delta'R^T \wedge N\delta'' \wedge \mathcal{A}\delta''M^T = 0 (\Leftrightarrow M\delta''\mathcal{A} \wedge \delta''N^TR = \text{diag}), \\ M\delta''A'^kL^{-1}B = N\delta''A'^kL^{-1}B = M\delta''\mathcal{A} \wedge d''RR^TA'^kL^{-1}B = N\delta''\mathcal{A} \wedge d''RR^TA'^kL^{-1}B = 0, \\ k = 0, 1, \dots, p-1, \quad A' = \dots \oplus J_p \oplus \dots \oplus J_1, \quad p \geq 0 \text{ being the SJ decomposition of } A'$$

(the last relations are relevant for IQWC and state that $M\delta''$, $N\delta''$, $M\delta''\mathcal{A} \wedge d''RR^T$, $N\delta''\mathcal{A} \wedge d''RR^T$ are not supported on the coordinates corresponding to the SJ block of $L^T f_1$ in A').

Further imposing $d' \wedge$ and $d'' \wedge$ on (3.11), using $d' \wedge d' = 0$, $d' \wedge d'' + d'' \wedge d' = 0$, $d'' \wedge d'' = 0$, the equations of (3.11) itself and (3.10) we finally get involution (that is no further conditions appear); note that for diagonal QWC (which form an open dense set in the set of (I)QWC) (3.11) is vacuous, so we have (3.10) already in involution.

While the differential system (3.9) together with its compatibility conditions (3.10)&(3.11) is interesting on its own, these compatibility conditions must further be extended with new conditions in order to describe multiply conjugate systems containing deformations of quadrics; in order to find these conditions we need to consider the space realization of solutions.

To extend x with the independent variables $u^{n+1}, \dots, u^{2n-1}$ (3.2), (3.3) and (3.6) will provide the needed information to obtain multiply conjugate systems containing deformations of QWC. For our problem we have $m = 2n - 1$, $d'x = [V_1 \dots V_{2n-1}]^T d'V$ with R solution of (3.10)&(3.11), V , Λ , V_j , Λ_j , $j = 1, \dots, 2n - 1$ solutions of the (homogeneous) differential part of (3.9) satisfying (3.8) and $\mathbf{a}_j = \lambda_j$, $j = 1, \dots, n$; from $x_{j+n+l} = (\log \lambda_j)_{n+l}x_j + (\log \mathbf{a}_{n+l})_j x_{n+l}$, $j = 1, \dots, n$, $l = 1, \dots, n - 1$ we get $x_{n+l} = ([V_1 \dots V_{2n-1}]^T N - [\Lambda_1 \dots \Lambda_{2n-1}]^T M)e_l \frac{1}{c_l}$, $d'(\log \mathbf{a}_{n+l}) = c_l e_l^T \mathcal{A}d'V$; from $(\mathbf{a}_{n+l})_{jk} = (\log \lambda_j)_k (\mathbf{a}_{n+l})_j + (\log \lambda_k)_j (\mathbf{a}_{n+l})_k$, $j, k = 1, \dots, n$, $j \neq k$, $l = 1, \dots, n - 1$ we get $\frac{1}{c_l} = e_l^T (\mathcal{A}V + f)$, $d'f = 0$, so $\mathbf{a}_{n+l} = e^{g_l} e_l^T (\mathcal{A}V + f)$, $d'g_l = 0$, $d''x = ([V_1 \dots V_{2n-1}]^T N\delta'' - [\Lambda_1 \dots \Lambda_{2n-1}]^T M\delta'')(\mathcal{A}V + f)$. We need $0 = d' \wedge d''x + d'' \wedge d'x = [V_1 \dots V_{2n-1}]^T R\delta' \wedge M\delta''A'f$ (so $f = 0$), $0 = d'' \wedge d''x = ([V_1 \dots V_{2n-1}]^T N\delta'' - [\Lambda_1 \dots \Lambda_{2n-1}]^T M\delta'')\mathcal{A}^2 \wedge (\delta''M^T\Lambda - \delta''N^TV)$, so we need

$$(3.12) \quad M\delta''\mathcal{A} \wedge \mathcal{A}\delta''M^T = N\delta''\mathcal{A} \wedge \mathcal{A}\delta''M^T = N\delta''\mathcal{A} \wedge \mathcal{A}\delta''N^T = 0.$$

Imposing $d' \wedge$ and $d'' \wedge$ conditions on (3.12) and using (3.10)&(3.11) and (3.12) themselves we don't get any new conditions; note also that (3.12) are vacuous for diagonal QWC. Now we need

$$\begin{aligned}
x_{n+l} n+p &= (\log \mathbf{a}_{n+l})_{n+p} x_{n+l} + (\log \mathbf{a}_{n+p})_{n+l} x_{n+p}, \quad l, p = 1, \dots, n-1, \quad l \neq p, \\
(\lambda_j)_{k} n+l &= (\log \lambda_k)_{n+l} (\lambda_j)_k + (\log \mathbf{a}_{n+l})_k (\lambda_j)_{n+l}, \quad j, k = 1, \dots, n, \quad j \neq k, \quad l = 1, \dots, n-1, \\
(\lambda_j)_{n+l} n+p &= (\log \mathbf{a}_{n+l})_{n+p} (\lambda_j)_{n+l} + (\log \mathbf{a}_{n+p})_{n+l} (\lambda_j)_{n+p}, \quad j = 1, \dots, n, \quad l, p = 1, \dots, n-1, \quad l \neq p, \\
(\mathbf{a}_{n+l})_j n+p &= (\log \lambda_j)_{n+p} (\mathbf{a}_{n+l})_j + (\log \mathbf{a}_{n+p})_j (\mathbf{a}_{n+l})_{n+p}, \quad j = 1, \dots, n, \quad l, p = 1, \dots, n-1, \quad l \neq p, \\
(\mathbf{a}_{n+l})_{n+p} n+q &= (\log \mathbf{a}_{n+p})_{n+q} (\mathbf{a}_{n+l})_{n+p} + (\log \mathbf{a}_{n+q})_{n+p} (\mathbf{a}_{n+l})_{n+q}, \quad l, p, q = 1, \dots, n-1 \text{ distinct};
\end{aligned}
\tag{3.13}$$

these are satisfied for diagonal QWC if $d'' g_l = (g_l)_{n+l} du^{n+l}$.

3.2. Quadrics with center.

4. THE BÄCKLUND TRANSFORMATION

4.1. (Isotropic) quadrics without center.

For the B transformation the space realization

$$(4.1) \quad x^1 = x^0 + [x_{v_1^0}^0 \dots x_{v_n^0}^0](\sqrt{R'_z} V_1 - V_0 + I_{1,n} L^{-1} C(z)) \subset \mathbb{C}^{2n-1}$$

of the leaf x^1 relative to the seed $x^0 \subset \mathbb{C}^{2n-1}$, $|d' x^0|^2 = |d' x_0^0|^2$, the algebraic transformation

$$\begin{aligned}
V_1 &= \sqrt{R'_z} V_0 - \sqrt{z} R_1 \Lambda_0 + I_{1,n} L^{-1} C(z), \quad \Lambda_1 = R_0^T (\sqrt{z} A' V_0 + \sqrt{R'_z} R_1 \Lambda_0 + \sqrt{z} I_{1,n} L^{-1} B), \\
(0, \sqrt{z}) &\leftrightarrow (1, -\sqrt{z})
\end{aligned}
\tag{4.2}$$

of solutions of (3.9) and the algebraic formula of the BPT

$$(4.3) \quad R_3 R_0^T = (D_2 - D_1 R_2 R_1^T)(D_2 R_2 R_1^T - D_1)^{-1}, \quad D_j := \sqrt{R'_{zj}} / \sqrt{z_j}, \quad j = 1, 2$$

remain valid, but the differential system subjacent to the B transformation (Ricatti equation)

$$(4.4) \quad -d' R_1 = R_1 \omega'_0 + R_1 \delta' R_0^T D R_1 - D R_0 \delta', \quad D := \sqrt{R'_z} / \sqrt{z}, \quad (0, \sqrt{z}) \leftrightarrow (1, -\sqrt{z})$$

in R_1 must be extended and then the BPT algebraic formula must satisfy this extension (for the third Möbius configuration the algebraic computations only suffice).

Applying d'' to (4.2) we get the full Ricatti equation

$$\begin{aligned}
d' R_1 &= -R_1 \omega'_0 - R_1 \delta' R_0^T D R_1 + D R_0 \delta', \\
(4.5) \quad d'' R_1 &= D^{-1} (d'' R_0 R_0^T D^2 + N_0 \delta'') D^{-1} R_1 + D^{-1} \delta'' M_0^T - R_1 M_0 \delta'' D^{-1} R_1
\end{aligned}$$

in R_1 and further

$$\begin{aligned}
M_1 \delta'' &= R_0^T (D R_1 M_0 \delta'' - N_0 \delta'') D^{-1}, \quad N_1 \delta'' = (A' R_1 M_0 \delta'' - R_1 M_0 \delta'' A' + D N_0 \delta'') D^{-1} \\
(4.6)
\end{aligned}$$

(thus we need $A' e_n \subset \mathbb{C} e_n$).

Imposing compatibility conditions $d' \wedge$, $d'' \wedge$ on (4.5) and using the equation itself we get $d' \wedge d' R_1 = R_1 (-d' \wedge \omega'_0 + \omega'_0 \wedge \omega'_0 - \delta' R_0^T A' R_0 \wedge \delta') - D R_0 (\delta' \wedge \omega'_0 - R_0^T d' R_0 \wedge \delta') + R_1 (\omega'_0 \wedge \delta' - \delta' \wedge R_0^T d' R_0) R_0^T D R_1$, $d' \wedge d'' R_1 + d'' \wedge d' R_1 = D^{-1} [d' \wedge (d'' R_0 R_0^T) - \delta'' M_0^T \wedge \delta' R_0^T - R_0 \delta' \wedge M_0 \delta''] D R_1 - R_1 [d'' \wedge \omega'_0 - M_0 \delta'' \wedge R_0 \delta' - \delta' R_0^T \wedge \delta'' M_0^T] - R_1 [d' \wedge (M_0 \delta'') - \omega'_0 \wedge M_0 \delta'' - \delta' R_0^T \wedge N_0 \delta''] D^{-1} R_1 + D^{-1} [d' \wedge (N_0 \delta'') - R_0 \delta' \wedge M_0 \delta'' A' + A' R_0 \delta' \wedge M_0 \delta''] D^{-1} R_1 + D^{-1} [d' \wedge (\delta'' M_0^T) - \delta'' M_0^T \wedge \omega'_0 + \delta'' N_0^T \wedge R_0 \delta'], \quad d'' \wedge d'' R_1 = D^{-1} [d'' \wedge (N_0 \delta'') - \delta'' M_0^T \wedge M_0 \delta'' - N_0 \delta'' \wedge (d'' R_0 R_0^T - D^{-2} \delta'' N_0^T) - d'' R_0 R_0^T \wedge N_0 \delta''] D^{-1} R_1 + D^{-1} [d'' \wedge (\delta'' M_0^T) - (d'' R_0 R_0^T + N_0 \delta'' D^{-2}) \wedge \delta'' M_0^T] - R_1 [d'' \wedge (M_0 \delta'') - M_0 \delta'' \wedge (d'' R_0 R_0^T - D^{-2} \delta'' N_0^T)] + R_1 M_0 \delta'' D^{-2} \wedge \delta'' M_0^T + (R_1 M_0 \delta'' - D^{-1} N_0 \delta'') D^{-2} \wedge (d'' R_0 R_0^T D^2 + N_0 \delta'' - D^2 d'' R_0 R_0^T + \delta'' N_0^T) D^{-1} R_1$; for diagonal QWC these are 0 from (3.10), so (4.5) is completely integrable and admits solution for any initial value R_1 . If the initial value is orthogonal, we would like the solution to remain orthogonal: $d(R_1 R_1^T - I_n) = -R_1 \delta' R_0^T D (R_1 R_1^T - I_n) - (R_1 R_1^T - I_n) D R_0 \delta' R_1^T - (R_1 R_1^T - I_n) D^{-1} (\delta'' M_0^T R_1^T + d'' R_0 R_0^T D^2 + N_0 \delta'') D^{-1}$, so $R_1 R_1^T - I_n$

is a solution of a linear differential equation and remains 0 if initially it was 0. The fact that R_1 is itself a solution of (3.10) follows from the symmetry $(0, \sqrt{z}) \leftrightarrow (1, -\sqrt{z})$ and the fact that $d \wedge dR_0 = 0$ (basically we use the converse of the proven results).

Therefore we only need to prove that R_3 given by (4.3) satisfies (4.5) for (R_0, z) replaced by (R_1, z_2) , (R_2, z_1) ; by symmetry it is enough to prove only one relation. Since $dR_1 = -R_1\omega'_0 - R_1\delta'R_0^T D_1 R_1 + D_1 R_0 \delta' + D_1^{-1}(d''R_0 R_0^T D_1^2 + N_0 \delta'')D_1^{-1} R_1 + D_1^{-1}\delta''M_0^T - R_1 M_0 \delta'' D_1^{-1} R_1$, $dR_2 = -R_2\omega'_0 - R_2\delta'R_0^T D_2 R_2 + D_2 R_0 \delta' + D_2^{-1}(d''R_0 R_0^T D_2^2 + N_0 \delta'')D_2^{-1} R_2 + D_2^{-1}\delta''M_0^T - R_2 M_0 \delta'' D_2^{-1} R_2$, we get $d(R_2 R_1^T) = [-R_2\omega'_0 - R_2\delta'R_0^T D_2 R_2 + D_2 R_0 \delta' + D_2^{-1}(d''R_0 R_0^T D_2^2 + N_0 \delta'')D_2^{-1} R_2 + D_2^{-1}\delta''M_0^T - R_2 M_0 \delta'' D_2^{-1} R_2]R_1^T + R_2[\omega'_0 R_1^T - R_1^T D_1 R_0 \delta' R_1^T + \delta' R_0^T D_1 + R_1^T D_1^{-1}(-D_1^2 d''R_0 R_0^T + \delta''N_0^T)D_1^{-1} + M_0 \delta'' D_1^{-1} - R_1^T D_1^{-1} \delta'' M_0^T R_1^T] = -(R_2 R_1^T)R_1 \delta' R_0^T (D_2 R_2 R_1^T - D_1) + (D_2 - R_2 R_1^T D_1)R_0 \delta' R_1^T - (R_2 R_1^T D_1^{-1} - D_2^{-1})\delta'' M_0^T R_1^T - R_2 M_0 \delta'' (D_2^{-1} R_2 R_1^T - D_1^{-1}) + D_2^{-1}(d''R_0 R_0^T D_2^2 + N_0 \delta'')D_2^{-1} R_2 R_1^T - R_2 R_1^T D_1^{-1}(d''R_0 R_0^T D_1^2 + N_0 \delta'')D_1^{-1}$. Thus if we prove the similar relation $d(R_3 R_0^T) = -(R_3 R_0^T)R_0 \delta' R_1^T (D_2 R_3 R_0^T + D_1) + (D_2 + R_3 R_0^T D_1)R_1 \delta' R_0^T + (R_3 R_0^T D_1^{-1} + D_2^{-1})\delta'' M_1^T R_0^T - R_3 M_1 \delta'' (D_2^{-1} R_3 R_0^T + D_1^{-1}) + D_2^{-1}(d''R_1 R_1^T D_2^2 + N_1 \delta'')D_2^{-1} R_3 R_0^T - R_3 R_0^T D_1^{-1}(d''R_1 R_1^T D_1^2 + N_1 \delta'')D_1^{-1}$, then since $dR_0 = -R_0\omega_1 + R_0\delta'R_1^T D_1 R_0 - D_1 R_1 \delta' + D_1^{-1}(d''R_1 R_1^T D_1^2 + N_1 \delta'')D_1^{-1} R_0 - D_1^{-1}\delta''M_1^T + R_0 M_1 \delta'' D_1^{-1} R_0$ we obtain what we want: $dR_3 = -R_3\omega_1 - R_3\delta'R_1^T D_2 R_3 + D_2 R_1 \delta' + D_2^{-1}(d''R_1 R_1^T D_2^2 + N_1 \delta'')D_2^{-1} R_3 + D_2^{-1}\delta''M_1^T - R_3 M_1 \delta'' D_2^{-1} R_3$. Differentiating (4.3) we get $d(R_3 R_0^T)(D_2 R_2 R_1^T - D_1) = -(R_3 R_0^T D_2 + D_1)d(R_2 R_1^T)$; thus we need to prove $[-(R_3 R_0^T)R_0 \delta' R_1^T (D_2 R_3 R_0^T + D_1) + (D_2 + R_3 R_0^T D_1)R_1 \delta' R_0^T + (R_3 R_0^T D_1^{-1} + D_2^{-1})\delta'' M_1^T R_0^T - R_3 M_1 \delta'' (D_2^{-1} R_3 R_0^T + D_1^{-1}) + D_2^{-1}(d''R_1 R_1^T D_2^2 + N_1 \delta'')D_2^{-1} R_3 R_0^T - R_3 R_0^T D_1^{-1}(d''R_1 R_1^T D_1^2 + N_1 \delta'')D_1^{-1}](D_2 R_2 R_1^T - D_1) = -(R_3 R_0^T D_2 + D_1)[-(R_2 R_1^T)R_1 \delta' R_0^T (D_2 R_2 R_1^T - D_1) + (D_2 - R_2 R_1^T D_1)R_0 \delta' R_1^T - (R_2 R_1^T D_1^{-1} - D_2^{-1})\delta'' M_0^T R_1^T - R_2 M_0 \delta'' (D_2^{-1} R_2 R_1^T - D_1^{-1}) + D_2^{-1}(d''R_0 R_0^T D_2^2 + N_0 \delta'')D_2^{-1} R_2 R_1^T - R_2 R_1^T D_1^{-1}(d''R_0 R_0^T D_1^2 + N_0 \delta'')D_1^{-1}]$. The terms containing $R_1 \delta' R_0^T$ become $D_2 + R_3 R_0^T D_1 = (R_3 R_0^T D_2 + D_1)R_2 R_1^T$ which follows directly from (4.3) and the terms containing $R_0 \delta' R_1^T$ become $R_0 \delta' R_1^T (D_2 R_3 R_0^T + D_1)(D_2 R_2 R_1^T - D_1) = (R_3 R_0^T)^T (R_3 R_0^T D_2 + D_1)(D_2 - R_2 R_1^T D_1)R_0 \delta' R_1^T$ which follows from $(D_2 R_3 R_0^T + D_1)(D_2 R_2 R_1^T - D_1) = D_2^2 - D_1^2 = (\frac{1}{z_2} - \frac{1}{z_1})I_n = (R_3 R_0^T)^T (R_3 R_0^T D_2 + D_1)(D_2 - R_2 R_1^T D_1)$.

Replacing $d''R_1 R_1^T$, $M_1 \delta''$, $N_1 \delta''$ from (4.5)&(4.6) and using $\delta''N_0^T = -A'd''R_0 R_0^T + d''R_0 R_0^T A' - N_0 \delta''$ the remaining terms split into ones containing $d''R_0 R_0^T$, $R_1 M_0 \delta''$, $\delta''M_0^T R_1^T$, $N_0 \delta''$ and they turn out to be as they should.

For the terms containing $\Omega := d''R_0 R_0^T$ we need $[D_2^{-1}D_1^{-1}\Omega D_1 D_2 R_3 R_0^T - R_3 R_0^T D_1^{-2}\Omega D_1^2 + (R_3 R_0^T D_1^{-1} + D_2^{-1})D_1^{-1}(A'\Omega - \Omega A')](D_2 R_2 R_1^T - D_1) + (R_3 R_0^T D_2 + D_1)(D_2^{-1}\Omega D_2 R_2 R_1^T - R_2 R_1^T D_1^{-1}\Omega D_1) = 0$; for the terms containing $\Omega := R_1 M_0 \delta''$ we need $[D_2^{-1}(-\Omega D_2 + (A'\Omega - \Omega A')D_2^{-1})D_1^{-1}R_3 R_0^T + R_3 R_0^T D_1^{-1}(\Omega - (A'\Omega - \Omega A')D_1^{-1}) - R_3 R_0^T D_1 \Omega D_1^{-1}(D_2^{-1}R_3 R_0^T - D_1^{-1})](D_2 R_2 R_1^T - D_1) - (R_3 R_0^T D_2 + D_1)R_2 R_1^T \Omega (D_2^{-1}R_2 R_1^T - D_1^{-1}) = 0$; for the terms containing $\Omega := D_2^{-1}D_1^{-1}\delta''M_0^T R_1^T$ we need $\Omega(D_2 R_3 R_0^T + D_1)(D_2 R_2 R_1^T - D_1) - (R_3 R_0^T D_2 + D_1)(R_2 R_1^T D_2 - D_1)\Omega = 0$; for the terms containing $\Omega := N_0 \delta''$ we need $[D_2^{-1}D_1^{-1}(\Omega D_2^2 + D_1^2 \Omega)D_2^{-1}D_1^{-1}R_3 R_0^T - R_3 R_0^T (D_1^{-2}\Omega + \Omega D_1^{-2}) + (R_3 R_0^T D_1^{-1} + D_2^{-1})D_1^{-1}\Omega + R_3 R_0^T \Omega D_1^{-1}(D_2^{-1}R_3 R_0^T - D_1^{-1})](D_2 R_2 R_1^T - D_1) + (R_3 R_0^T D_2 + D_1)[D_2^{-1}\Omega D_2^{-1}R_2 R_1^T - R_2 R_1^T D_1^{-1}\Omega D_1^{-1}] = 0$.

4.2. Quadrics with center.

REFERENCES

- [1] Berger, E., Bryant, R. L., Griffiths, P. A. *The Gauß equations and rigidity of isometric embeddings*, Duke Math. J., **50** (1983), 803-892.
- [2] L. Bianchi *Sur la déformation des quadriques*, Comptes rendus de l'Académie, **142**, (1906), 562-564; and **143**, (1906) 633-635.
- [3] L. Bianchi *Lezioni Di Geometria Differenziale, Teoria delle Trasformazioni delle Superficie applicabili sulle quadriche*, Vol **3**, Enrico Spoerri Libraio-Editore, Pisa (1909).
- [4] L. Bianchi *Lezioni Di Geometria Differenziale*, Vol **1-4**, Nicola Zanichelli Editore, Bologna (1922-27).
- [5] L. Bianchi *Opere*, Vol **1-11**, a cura dell'Unione Matematica Italiana e col contributo del Consiglio Nazionale Delle Ricerche, Edizioni Cremonese (1952-59).

- [6] A. I. Bobenko, U. Pinkall *Discrete surfaces with constant negative Gaussian curvature and the Hirota equation*, J. Diff. Geom., **43** (1996), no. 3, 527-611.
- [7] E. Cartan *Sur les variétés de courbure constante d'un espace euclidien ou non-euclidien*, Bull. Soc. Math. France, **47** (1919), 125-160; and **48** (1920), 132-208.
- [8] G. Darboux *Leçons Sur La Théorie Générale Des Surfaces*, Vol **1-4**, Gauthier-Villars, Paris (1894-1917).
- [9] G. Darboux *Leçons Sur Les Systèmes Orthogonaux Et Les Coordonnées Curvilignes*, Gauthier-Villars, Paris (1910).
- [10] G. M. Green *Projective Differential Geometry of Triple Systems of Surfaces*, Press of The New Era Printing Company, Lancaster, PA (1913).
- [11] K. -M. Peterson *Sur la déformation des surfaces du second ordre*, Annales de la faculté des sciences de Toulouse 2^e série, tome 7, **1** (1905), 69-107.
- [12] K. Tenenblat and C.-L. Terng *Bäcklund's theorem for n-dimensional sub-manifolds of $\mathbb{R}^{2n-1}$* , Ann. of Math., **111** (1980), 477-490.
- [13] C.-L. Terng *A higher dimension generalization of the Sine-Gordon equation and its soliton theory*, Ann. of Math., **111** (1980), 491-510.
- [14] C.-L. Terng, K. Uhlenbeck *Poisson Actions and Scattering Theory for Integrable Systems*, arXiv:dg-ga/9707004 v1 7 Jul 1997.

FACULTY OF MATHEMATICS AND INFORMATICS, UNIVERSITY OF BUCHAREST, 14 ACADEMIEI STR., 010014, BUCHAREST, ROMANIA

E-mail address: `dinca@gta.math.unibuc.ro`